

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/131072/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Basak, Shyam, Alvarez-Montoya, Ana, Winfrey, Laura, Melen, Rebecca L. ORCID: <https://orcid.org/0000-0003-3142-2831>, Morrill, Louis C. ORCID: <https://orcid.org/0000-0002-6453-7531> and Pulis, Alexander P. 2020. B(C6F5)3-catalyzed direct C3 alkylation of indoles and oxindoles. ACS Catalysis 10 (8) , pp. 4835-4840. 10.1021/acscatal.0c01141 file

Publishers page: <http://dx.doi.org/10.1021/acscatal.0c01141>
<<http://dx.doi.org/10.1021/acscatal.0c01141>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



B(C₆F₅)₃-Catalyzed Direct C3 Alkylation of Indoles and Oxindoles

Shyam Basak, Ana Alvarez-Montoya, Laura Winfrey, Rebecca L. Melen,* Louis C. Morrill,* and Alexander P. Pulis*

Cite This: *ACS Catal.* 2020, 10, 4835–4840

Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: The direct C3 alkylation of indoles and oxindoles is a challenging transformation, and only a few direct methods exist. Utilizing the underexplored ability of triaryl boranes to mediate the heterolytic cleavage of α -nitrogen C–H bonds in amines, we have developed a catalytic approach for the direct C3 alkylation of a wide range of indoles and oxindoles using amine-based alkylating agents. We also employed this borane-catalyzed strategy in an alkylation–ring opening cascade.



KEYWORDS: catalysis, boranes, *N*-heterocycles, alkylation, indoles, oxindoles

Indoles and oxindoles are prevalent motifs in biologically active molecules.¹ Classic indole syntheses involve ring construction.² Another approach involves the functionalization of the readily accessible heterocycle core; yet, the direct and selective C3 alkylation of indoles and oxindoles is a surprisingly challenging transformation as the reaction with simple alkyl halides is often not synthetically useful.^{2,3} For example, with methyl iodide, 1,2-dimethylindole and 1-methylindole are unreactive,⁴ 2-methylindole results in mixtures of *N*- and *C*-methylation,⁵ and oxindoles undergo dialkylation at C3.³ The installation of a methyl group is a worthwhile endeavor, considering the interest of medicinal chemists in the “magic methyl effect”;⁶ yet only a few methods exist for the direct C3 methylation of indoles and oxindoles (Scheme 1a). Direct C3 methylation is possible with CO₂/H₂ and a ruthenium catalyst (e.g., for 1,2-dimethylindole and 2-methylindole),⁷ and with borrowing hydrogen methods with methanol (e.g., for 2-methylindole⁸ and 1-phenyl oxindole).^{8a,9} The direct methylation of 1-methylindole is currently unknown.⁴

Because of their intrinsic Lewis acidity, borane catalysts have found numerous applications in synthesis and are traditionally used to activate polarized bonds.¹⁰ Triaryl boranes can also activate unpolarized bonds, such as H–H¹¹ and Si–H bonds.¹² In a similar vein, we considered if boranes could also be used to cleave C(sp³)–H bonds heterolytically¹³ and unveil new approaches to challenging transformations. Related to this, we were intrigued by a report by Santini that described the heterolytic cleavage of an α -nitrogen C(sp³)–H bond during the stoichiometric reaction of dimethyl aniline and B(C₆F₅)₃ to form an iminium borohydride ion pair (Scheme 1b).¹⁴ B(C₆F₅)₃-mediated α -N C(sp³)–H bond cleavage¹⁵ was

unrecognized as a synthetic strategy for almost a decade until Stephan and co-workers reported its use in the transfer hydrogenation of imines.¹⁶ Subsequently, Grimme and Paradies,^{17a} Kanai,^{17b} and Zhang^{17c} disclosed methods for the dehydrogenation of *N*-heterocycles. A major breakthrough came when Erker reported the use of this unusual reactivity in C–C bond-forming reactions where stoichiometric B(C₆F₅)₃ was used to generate iminium ions for Mannich-type processes.¹⁸ Wasa greatly advanced the strategy by reporting the catalytic use of B(C₆F₅)₃ in an asymmetric Mannich process.¹⁹ The iminium ions generated have also been used in electrocyclizations,²⁰ and in the β -functionalization of amines.^{21,22} However, the use of this reactivity in catalytic C–C bond-forming reactions remains rare.^{19,20} Inspired by these reports and borrowing hydrogen alkylation reactions,²³ we have applied this underutilized reactivity in challenging alkylation processes.

Here, we have developed a new strategy for the direct C3 methylation of indoles and oxindoles (Scheme 1c). The process utilizes a B(C₆F₅)₃-mediated α -N C(sp³)–H bond cleavage events to activate readily available amine-based alkylating agents. Using this borane-catalyzed method, common undesired reactions, such as the *N*-methylation of

Received: March 9, 2020

Revised: April 2, 2020

Published: April 9, 2020




Scheme 1. B(C₆F₅)₃-Catalyzed α -N C(sp³)-H Bond Cleavage Used in the Methylation of Indoles and Oxindoles


a) Synthetic challenge: Direct C3 methylation of indoles and oxindoles

Current methods for direct C3 methylation:

CO₂ (20 atm), H₂, [Ru] cat.⁷ only (TM = Pt, Ir, Co, Fe)
 ref. 7 and CH₃OH, [TM] cat.⁸ only
 CH₃OH, [Fe]^{8a} or [Pd]⁹ cat. only




b) B(C₆F₅)₃-mediated heterolytic C-H bond cleavage



Previously reported uses:

- Transfer hydrogenation/dehydrogenation
- Mannich type reactions
- Iminium electrocyclozation
- Enamine formation

c) B(C₆F₅)₃-catalyzed direct C3 alkylation of indoles and oxindoles (this work)




Beyond methylation: other alkylations including alkylation/ring opening cascades


indoles, the formation of 3,3'-bisindolylmethanes, and the dialkylation of oxindoles, are not observed. In addition, the substrate scope is broad and encompasses 1-, 2-, and 1,2-substituted indoles, as well as other challenging alkylations, including a novel alkylation-ring opening cascade.

We began by investigating various aniline derivatives as methylating agents in the borane-catalyzed methylation of 1,2-dimethyl indole (**1a**) (Scheme 2). Generally, we discovered

Scheme 2. B(C₆F₅)₃-Catalyzed Methylation of Indole **1a** with Various Alkylating Agents^a



methylating agents:



^aReactions were performed using 0.2 mmol of **1a**. Yields were determined after ¹H NMR spectrum analysis of the crude reaction mixture with an internal standard.

that a variety of aryl and diaryl amines were effective in methylating **1a** using B(C₆F₅)₃ (10 mol %).²⁴ Electron-rich diaryl methyl amines, such as **4a** and **6a**, were determined to be optimal and allowed the formation of **2a** in quantitative yields at ambient temperature.

We surveyed the scope of the B(C₆F₅)₃-catalyzed methylation of various 1,2-, 1-, and 2-substituted indoles and oxindoles and found that the reaction broadly tolerated a range of functional groups and substitution patterns (Scheme


3). Notably, the direct methylation of 1-methylindole (**1f**), which is a transformation that was previously absent from the literature,⁴ was successfully accomplished in high isolated yield (**2f**, 75%) using the B(C₆F₅)₃-catalyzed approach with methylating agent **6a**.²⁵ 2-Substituted indoles (i.e., NH indoles, cf. **2l–2s**) were efficiently methylated when 2,2,6,6-tetramethylpiperidine (TMP, 10 mol %) was used with alkylating agent **6a** and B(C₆F₅)₃ (10 mol %).²⁶ Importantly, *N*-methylation was not observed with NH-bearing indoles. In contrast, *N*-alkylation, or mixtures of *N*- and *C*-alkylation, typically result when NH indoles are treated with methyl iodide under basic conditions.⁵ The successful reaction of 1- (cf. **2f–2k**) and 2-substituted indoles (cf. **2l–2s**) was surprising, given that B(C₆F₅)₃ has been reported to react readily with these classes of heterocycle to produce zwitterionic species.²⁷ 3,3'-Bisindolylmethanes, which are a common product formed in the reaction of formaldehyde or iminium electrophiles with indoles, were not observed.²⁸

Oxindoles (**8a–8q**) were successfully employed in the B(C₆F₅)₃-catalyzed methylation to give products **9a–9q**. In this class of heterocycle, 1,2,2,6,6-pentamethylpiperidine (PMP, **13**) was used as the alkylating agent and higher temperatures were required. Crucially, C3 dimethylation was not observed. Therefore, the borane-catalyzed process complements traditional alkylating agents: C3 dialkylation typically occurs when oxindoles are treated with methyl iodide under basic conditions.³


The methylation of 6-methylindole (cf. **2n**) and unsubstituted oxindole (cf. **9n**) occurred in low yield, presumably because of competitive coordination of N or O to the B(C₆F₅)₃ catalyst. Otherwise, across the different classes of substrates, the process tolerated a range of functional groups and substituents, such as OCH₃ (**2c**, **2s**, **9i**, **9k**), F (**2o**, **9d**), Cl (**2d**, **2p**, **9e**), Br (**2q**, **9f**), CF₃ (**9m**), NO₂ (**2e**, **9j**), CO₂Me (**9c**), and other carbonyl derivatives (**9o**, **9p**), which contrasts the dogma sometimes associated with B(C₆F₅)₃-mediated processes.²⁹ We also performed the B(C₆F₅)₃-catalyzed methylation of 1,2-dimethylindole (**1a**) on a preparative scale, producing 1.3 g of 1,2,3-trimethylindole (**2a**) in 83% yield.³⁰

In addition, we briefly explored other challenging alkylation reactions using the B(C₆F₅)₃-catalyzed method and discovered that 1,2-dimethylindole (**1a**) was successfully ethylated (**10a**), decylated (**11a**) and benzylated (**12a**), at C3 using the ethyl- (**6b**), decyl- (**6c**), or benzyl- (**4b**)³¹ diaryl amines, respectively.³²

The borane catalyst, B(C₆F₅)₃, is a commercially available white powder that forms a water adduct, H₂O·B(C₆F₅)₃, when exposed to moisture in air and is therefore routinely handled in an inert atmosphere.³³ Inspired by related methods,³⁴ we developed a procedure where B(C₆F₅)₃ can be used as received from the supplier and weighed in air on the open bench, and the reaction performed using standard Schlenk line techniques (Scheme 4). Thus, H₂O·B(C₆F₅)₃ (10 mol %) was dissolved in the desired solvents (as received from the supplier) and treated with triethyl silane (20 mol %). The resultant solution contains active B(C₆F₅)₃ and O(SiEt₃)₂ that can be used directly in the alkylation of indoles and oxindoles to provide methylated indoles (**2a**, **2f**, and **2l**), benzylated indole (**12a**), and methylated oxindole (**9a**)³⁵ in good yields. Therefore, this shows that access to specialized equipment (such as a dry glovebox), a separate purification of commercially available

Scheme 3. Substrate Scope in the $B(C_6F_5)_3$ -Catalyzed Alkylation of Indoles and Oxindoles*


*Reactions were performed using 0.5 mmol of **1** or **8** except conditions **g** and **h**, where 0.2 mmol of **1a** was used. Yields are isolated. Yields in parentheses determined after 1H NMR spectrum analysis of the crude reaction mixture with an internal standard. ^a $B(C_6F_5)_3$ (10 mol %), **6a** ($R = CH_3$, 1.2 equiv), 25 °C, DCE, 16 h. ^b $B(C_6F_5)_3$ (10 mol %), **6a** ($R = CH_3$, 1.2 equiv), 95 °C, DCE, 16 h. ^c $B(C_6F_5)_3$ (20 mol %), **6a** ($R = CH_3$, 1.2 equiv), 95 °C, DCE, 8 h. ^d $B(C_6F_5)_3$ (10 mol %), **6a** ($R = CH_3$, 1.2 equiv), TMP (10 mol %), 110 °C, toluene, 16 h. ^e $B(C_6F_5)_3$ (10 mol %), **13** (PMP, 2 equiv), 150 °C, xylenes, 16 h. ^fCombined yield of tautomers. ^g $B(C_6F_5)_3$ (10 mol %), **6b** ($R = Et$) or **6c** ($R = (CH_2)_9CH_3$) (1.2 equiv), 95 °C, DCE, 24 h. ^h $B(C_6F_5)_3$ (20 mol %), **4b** ($R = Bn$, 2 equiv), 150 °C, xylenes, 24 h.

Scheme 4. Use of $H_2O \cdot B(C_6F_5)_3$ in the Borane-Catalyzed Alkylation of Indoles and Oxindoles

^a**6a**, 25 °C, DCE, 16 h. ^b**6a**, $B(C_6F_5)_3$ (20 mol %), Et_3SiH (40 mol %), 95 °C, DCE, 8 h. ^c**6a**, TMP (10 mol %), 110 °C, toluene, 16 h. ^d**4b**, 150 °C, *p*-xylene, 24 h. ^ePMP (**13**) (2 equiv), 150 °C, *p*-xylene, 16 h.

$B(C_6F_5)_3$, and rigorously anhydrous solvent is not required in the $B(C_6F_5)_3$ -catalyzed alkylation.

Beyond methylation and alkylation, we also explored the $B(C_6F_5)_3$ -catalyzed alkylation strategy in a novel alkylation-ring opening cascade process for the generation of functionalized indoles **15** (Scheme 5). Product **15** contains a 4-(3-indolyl)butylamine motif that is found in several serotonergic/dopaminergic drug molecules, such as vilazodone, roxindole, siramesine, and carmoxirole.³⁶ Upon reaction of *N*-aryl


Scheme 5. $B(C_6F_5)_3$ -Catalyzed Alkylation-Ring Opening Cascade*

*Standard conditions: $H_2O \cdot B(C_6F_5)_3$ (10 mol %), Et_3SiH (20 mol %), **14** (1 equiv), **1** (2.2 equiv), 1,2- $Cl_2C_6H_4$, 110 °C, 20–24 h. ^aDCE, 85 °C. ^bToluene.

pyrrolidines **14**,³⁷ indoles **1** and $B(C_6F_5)_3$ catalyst, a variety of 4-(3-indolyl)butylamines **15** were formed in good yields.³⁸

In order to probe the mechanism and provide direct access to deuterated methyl groups at C3 of indoles, we used deuterated methylating agent **6a-d₃** in the $B(C_6F_5)_3$ -catalyzed methylation of indoles **1a** and **1l** under previously optimized conditions (Scheme 6a). Deuterated C3 methylindoles **2a-d₃** and **2l-d₃** were formed in high yield in both cases.³⁹

Scheme 6. Mechanistic Studies and Proposed Catalytic Cycle*



*Yields are isolated and %D incorporation was determined after 1H NMR spectrum analysis of the purified compounds. ^aDCE, 25 °C, 16 h. ^bTMP (10 mol %), toluene, 110 °C, 16 h.

Based on these results and literature precedent, we propose the following catalytic cycle for the $B(C_6F_5)_3$ -catalyzed alkylation of indoles and oxindoles (Scheme 6b). The borane-catalyst mediates heterolytic cleavage, via hydride abstraction, of the α -N C(sp³)-H bond in the amine-based alkylating agents (**3–7**, **13**, **14**) forming iminium-borohydride ion pairs **16** (Scheme 6b, step (i)). Analogous ion pairs have been observed by Santini and co-workers using NMR spectroscopy (cf. Scheme 1A).¹⁴ The electrophilic iminium **16** is trapped with an indole **1** (or oxindole **8**), forging a new C–C bond (step (ii)) in an analogous fashion to the Mannich reaction. Proton transfers enable the ion pair **17** to eliminate the amine **18** (which can be recovered from the reaction) via an E1_{CB}-type mechanism (step (iii)).⁴⁰ The α,β -unsaturated iminium-based ion pair **19** is reduced by the borohydride

counterion, producing the alkylated indoles **2** (and oxindoles **9**) and regenerating the borane-catalyst (step (iv)). In the boron-catalyzed alkylation/ring opening cascade process (cf. Scheme 5), the cyclic nature of the iminium **20** enables the amino fragment to be retained in product **15** after elimination (Scheme 6c).

In summary, we have developed a new approach to the direct C3 alkylation of indoles and oxindoles. Using a $B(C_6F_5)_3$ catalyst and amine-derived alkylating agents, we exploit the underexplored ability of boranes to cleave heterolytically α -N C(sp³)-H bonds in a catalytic C–C bond-forming reaction. This method provides a metal-free and complementary approach to the few existing methods for the direct C3 alkylation of indoles. Unlike other procedures, this $B(C_6F_5)_3$ -catalyzed methodology encompasses several classes of indole, including 1-, 2-, and 1,2-substituted indoles, and allows previously unreported direct methylations. The reaction displays broad scope and exceptional chemoselectivity, avoiding *N*-methylation and formation of 3,3'-bisindolyl-methanes in indole substrates, and dialkylation in oxindoles. Other alkylations are also reported, including a novel alkylation-ring opening cascade process to generate privileged 4-(3-indolyl)butylamines from *N*-aryl pyrrolidines.

■ ASSOCIATED CONTENT

Supporting Information

Information about the data that underpins the results presented in this article, including how to access them, can be found in the Cardiff University data catalogue at <http://doi.org/10.17035/d.2020.0104936560> or from the lead authors upon request. The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscatal.0c01141>.

Experimental procedures and spectroscopic data (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

Rebecca L. Melen – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom; orcid.org/0000-0003-3142-2831; Email: MelenR@cardiff.ac.uk

Louis C. Morrill – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom; orcid.org/0000-0002-6453-7531; Email: MorrillLC@cardiff.ac.uk

Alexander P. Pulis – School of Chemistry, University of Leicester, Leicester LE1 7RH, United Kingdom; orcid.org/0000-0003-1754-527X; Email: a.pulis@leicester.ac.uk

Authors

Shyam Basak – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom

Ana Alvarez-Montoya – School of Chemistry, University of Leicester, Leicester LE1 7RH, United Kingdom

Laura Winfrey – School of Chemistry, University of Leicester, Leicester LE1 7RH, United Kingdom

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acscatal.0c01141>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

S.B., R.L.M., and L.C.M. gratefully acknowledge the School of Chemistry, Cardiff University for generous support and the Leverhulme Trust for a Research Project Grant (No. RPG-2015-361). R.L.M. would like to acknowledge the EPSRC for a research fellowship (No. EP/R026912/1). A.A.-M., L.W., and A.P.P. thank the School of Chemistry, University of Leicester, for their generous support and the Royal Society for a Research Grant (No. RGS\R1\191082).

■ REFERENCES

- (1) *Key Heterocycle Cores for Designing Multitargeting Molecules*; Silakari, O., Ed.; Elsevier, 2018; p 1.
- (2) Joule, J. A. Indole and Its Derivatives. In *Science of Synthesis: Knowledge Updates*, Vol. 2; Thomas, E. J., Ed.; Thieme Publishers Stuttgart, Germany, 2010; p 1.
- (3) Monoalkylation of oxindoles is often only achieved via indirect methods, for examples, see: (a) Bergman, J. Oxindoles. In *Advances in Heterocyclic Chemistry*, Vol. 117; Scriven, E. F. V.; Ramsden, C. A., Eds.; Elsevier, 2015; pp 1–81 (DOI: 10.1016/bs.aihch.2015.08.001). For examples of failed selective monoalkylation with methyl iodide, see: (b) Li, W.-H.; Li, C.; Dong, L. Highly Efficient Rhodium-Catalyzed Oxindole-Directed Oxidative Heck-Type Reaction of *N*-Aryloxindoles with Alkenes. *Asian J. Org. Chem.* **2018**, 7, 2448–5807. (c) Harrowven, D. C.; Stenning, K. J.; Whiting, S.; Thompson, T.; Walton, R. CH Activation and CH₂ Double Activation of Indolines by Radical Translocation: Understanding the Chemistry of the Indolinyl Radical. *Org. Biomol. Chem.* **2011**, 9, 4882–4885.
- (4) Scifinder and Reaxys database reaction searches (performed March 9, 2020).
- (5) After metalation, the reaction of 2-methyl indole with methyl iodide favors *N*-selectivity (in the case of sodium) or is unselective (in the case of magnesium, *N*-, mono *C*-, and bis *C*-alkylation results). For examples, see: (a) Ciszewski, L. W.; Durka, K.; Gryko, D. Photocatalytic Alkylation of Pyrroles and Indoles with α -Diazo Esters. *Org. Lett.* **2019**, 21, 7028–7032. (b) Rodríguez, J. G.; Urrutia, A. Acylation, Cyanoethylation and Alkylation of Methyl and Phenyl Indolylmagnesium Salts: Influence Of the Substituents on the *C*- and *N*-Reaction Products. *J. Heterocycl. Chem.* **1999**, 36, 129–135.
- (6) Schönherr, H.; Cernak, T. Profound Methyl Effects in Drug Discovery and a Call for New C–H Methylation Reactions. *Angew. Chem., Int. Ed.* **2013**, 52, 12256–12267.
- (7) Li, Y.; Yan, T.; Junge, K.; Beller, M. Catalytic Methylation of C–H Bonds Using CO₂ and H₂. *Angew. Chem., Int. Ed.* **2014**, 53, 10476–10480.
- (8) (a) Polidano, K.; Allen, B. D. W.; Williams, J. M. J.; Morrill, L. C. Iron-Catalyzed Methylation Using the Borrowing Hydrogen Approach. *ACS Catal.* **2018**, 8, 6440–6445. (b) Siddiki, S. M. A. H.; Touchy, A. S.; Jamil, M. A. R.; Toyao, T.; Shimizu, K. C-Methylation of Alcohols, Ketones, and Indoles with Methanol Using Heterogeneous Platinum Catalysts. *ACS Catal.* **2018**, 8, 3091–3103. (c) Seck, C.; Mbaye, M. D.; Gaillard, S.; Renaud, J.-L. Bifunctional Iron Complexes Catalyzed Alkylation of Indoles. *Adv. Synth. Catal.* **2018**, 360, 4640–4645. (d) Liu, Z.; Yang, Z.; Yu, X.; Zhang, H.; Yu, B.; Zhao, Y.; Liu, Z. Methylation of C(sp³)-H/C(sp²)-H Bonds with Methanol Catalysed by Cobalt System. *Org. Lett.* **2017**, 19, 5228–5231. (e) Chen, S.-J.; Lu, G.-P.; Cai, C. Iridium-Catalyzed Methylation of Indoles and Pyrroles using Methanol as Feedstock. *RSC Adv.* **2015**, 5, 70329–70332.
- (9) Mamidala, R.; Biswal, P.; Subramani, M. S.; Samser, S.; Venkatasubbaiah, K. Palladacycle-Phosphine Catalyzed Methylation of Amines and Ketones Using Methanol. *J. Org. Chem.* **2019**, 84, 10472–10480.
- (10) Ishihara, K. Achiral B(III) Lewis Acids. In *Lewis Acids in Organic Synthesis*; Yamamoto, H., Ed.; Wiley–VCH Verlag GmbH, 2000; p 1.
- (11) For recent reviews, see: (a) Stephan, D. W. Frustrated Lewis Pairs. *J. Am. Chem. Soc.* **2015**, 137, 10018–10032. (b) Stephan, D. W.; Erker, G. Frustrated Lewis Pair Chemistry: Development and Perspectives. *Angew. Chem., Int. Ed.* **2015**, 54, 6400–6441. (c) Stephan, D. W. The broadening Reach of Frustrated Lewis Pair Chemistry. *Science* **2016**, 354, aaf7229. (d) Pulis, A. P.; Winfrey, L.; Alvarez-Montoya, A. Uncovering the Synthetic Potential of Frustrated Lewis Pairs. In *Reference Module in Chemistry, Molecular Sciences and Chemical Engineering*; Elsevier, 2019; p 1.
- (12) For seminal reports, see: (a) Parks, D. J.; Piers, W. E. Tris(pentafluorophenyl)boron-Catalyzed Hydrosilylation of Aromatic Aldehydes, Ketones, and Esters. *J. Am. Chem. Soc.* **1996**, 118, 9440–9441. (b) Houghton, A. Y.; Hurmalainen, J.; Mansikkamäki, A.; Piers, W. E.; Tuononen, H. M. Direct Observation of a Borane–Silane Complex Involved in Frustrated Lewis-Pair-Mediated Hydrosilylations. *Nat. Chem.* **2014**, 6, 983–988. (c) Rendler, S.; Oestreich, M. Conclusive Evidence for an SN₂-Si Mechanism in the B(C₆F₅)₃-Catalyzed Hydrosilylation of Carbonyl Compounds: Implications for the Related Hydrogenation. *Angew. Chem., Int. Ed.* **2008**, 47, 5997–6000. For reviews, see ref 11d and (d) Oestreich, M.; Hermeke, J.; Mohr, J. A Unified Survey of Si–H and H–H Bond Activation Catalysed by Electron-Deficient Boranes. *Chem. Soc. Rev.* **2015**, 44, 2202–2220.
- (13) Systems consisting of boranes and Lewis bases react with C(sp³)-H bonds via deprotonation. For a review, see ref 11d. For examples of C(sp²)-H bond activation, see: (a) Légraré, M.-A.; Courtemanche, M.-A.; Rochette, É.; Fontaine, F.-G. Metal-Free Catalytic C–H bond Activation and Borylation of Heteroarenes. *Science* **2015**, 349, 513–516. (b) Chernichenko, K. C.; Lindqvist, M.; Kotai, B.; Nieger, M.; Sorochkina, K.; Papai, I.; Repo, T. Metal-Free sp²-C–H Borylation as a Common Reactivity Pattern of Frustrated 2-Aminophenylboranes. *J. Am. Chem. Soc.* **2016**, 138, 4860–4868.
- (14) Millot, N.; Santini, C. C.; Fenet, B.; Basset, J. M. Formation and Characterization of Zwitterionic Stereoisomers from the Reaction of B(C₆F₅)₃ and NEt₂Ph: (*E*)- and (*Z*)-[EtPhN⁺=CHCH₂-B[−](C₆F₅)₃]. *Eur. J. Inorg. Chem.* **2002**, 2002, 3328–3335.
- (15) Current methods to activate α -N C(sp³)-H bonds typically use oxidative conditions and precious-metal catalysts. For reviews, see: (a) Bruneau, C.; Achard, M. Alkane-Related C–H Bond Activation and Functionalization of Aliphatic Amines. In *Alkane Functionalization*; Pombeiro, A. K. L.; Guedes da Silva, M. F. C., Eds.; John Wiley & Sons, 2018; p 1. (b) Gini, A.; Brandhofer, T.; Mancheño, O. G. Recent Progress in Mild Csp³-H bond Dehydrogenative or (Mono-) Oxidative Functionalization. *Org. Biomol. Chem.* **2017**, 15, 1294–1312. (c) Cheng, M.-X.; Yang, S. D. Recent Advances in the Enantioselective Oxidative α -C–H Functionalization of Amines. *Synlett* **2017**, 28, 159–174.
- (16) Farrell, J. M.; Heiden, Z. M.; Stephan, D. W. Metal-Free Transfer Hydrogenation Catalysis by B(C₆F₅)₃. *Organometallics* **2011**, 30, 4497–4500.
- (17) (a) Maier, A. G. G.; Tussing, S.; Schneider, T.; Flörke, U.; Qu, Z.-W.; Grimme, S.; Paradies, J. Frustrated Lewis Pair Catalyzed Dehydrogenative Oxidation of Indolines and Other Heterocycles. *Angew. Chem., Int. Ed.* **2016**, 55, 12219–12223. (b) Kojima, M.; Kanai, M. Tris(pentafluorophenyl)borane-Catalyzed Acceptorless Dehydrogenation of *N*-Heterocycles. *Angew. Chem., Int. Ed.* **2016**, 55, 12224–12227. (c) Han, Y.; Zhang, S.; He, J.; Zhang, Y. B(C₆F₅)₃-Catalyzed (Convergent) Disproportionation Reaction of Indoles. *J. Am. Chem. Soc.* **2017**, 139, 7399–7407. (d) Zhang, S.; Han, Y.; He, J.; Zhang, Y. B(C₆F₅)₃-Catalyzed C3-Selective C–H Borylation of Indoles: Synthesis, Intermediates, and Reaction Mechanism. *J. Org. Chem.* **2018**, 83, 1377–1386.
- (18) Chen, G.-Q.; Kehr, G.; Daniliuc, C. G.; Bursch, M.; Grimme, S.; Erker, G. Intermolecular Redox-Neutral Amine C–H Functionalization Induced by the Strong Boron Lewis Acid B(C₆F₅)₃ in the Frustrated Lewis Pair Regime. *Chem. - Eur. J.* **2017**, 23, 4723–4729.
- (19) (a) Shang, M.; Chan, J. Z.; Cao, M.; Chang, Y.; Wang, Q.; Cook, B.; Torker, S.; Wasa, M. C–H Functionalization of Amines via Alkene-Derived Nucleophiles through Cooperative Action of Chiral

and Achiral Lewis Acid Catalysts: Applications in Enantioselective Synthesis. *J. Am. Chem. Soc.* **2018**, *140*, 10593–10601. (b) Chan, J. Z.; Chang, Y.; Wasa, M. $B(C_6F_5)_3$ -Catalyzed C–H Alkylation of N-Alkylamines Using Silicon Enolates without External Oxidant. *Org. Lett.* **2019**, *21*, 984–988.

(20) (a) Maier, A. F. G.; Tussing, S.; Zhu, H.; Wicker, G.; Tzvetkova, P.; Flörke, U.; Daniliuc, C. G.; Grimme, S.; Paradies, J. Borane-Catalyzed Synthesis of Quinolines Bearing Tetrasubstituted Stereocenters by Hydride Abstraction-Induced Electrocyclization. *Chem. - Eur. J.* **2018**, *24*, 16287–16291. (b) Tian, J.-J.; Zeng, N.-N.; Liu, N.; Tu, X.-S.; Wang, X. C. Intramolecular Cyclizations of Vinyl-Substituted *N,N*-Dialkyl Arylamines Enabled by Borane-Assisted Hydride Transfer. *ACS Catal.* **2019**, *9*, 295–300.

(21) (a) Zhang, J.; Park, S.; Chang, S. Catalytic Access to Bridged Sila-N-Heterocycles from Piperidines via Cascade sp^3 and sp^2 C–Si Bond Formation. *J. Am. Chem. Soc.* **2018**, *140*, 13209–13213. (b) Li, R.; Chen, Y.; Jiang, K.; Wang, F.; Lu, C.; Nie, J.; Chen, Z.; Yang, G.; Chen, Y.-C.; Zhao, Y.; Ma, C. $B(C_6F_5)_3$ -Catalyzed Redox-Neutral β -Alkylation of Tertiary Amines Using *p*-Quinone Methides via Borrowing Hydrogen. *Chem. Commun.* **2019**, *55*, 1217–1220. (c) Chang, Y.; Yesilcimen, A.; Cao, M.; Zhang, Y.; Zhang, B.; Chan, J. Z.; Wasa, M. Catalytic Deuterium Incorporation within Metabolically Stable β -Amino C–H Bonds of Drug Molecules. *J. Am. Chem. Soc.* **2019**, *141*, 14570–14575.

(22) Hydride abstraction from dienes and related compounds is known. For selected examples, see: (a) Simonneau, A.; Oestreich, M. Formal SiH_4 Chemistry Using Stable and Easy-To-Handle Surrogates. *Nat. Chem.* **2015**, *7*, 816–822. (b) Khan, I.; Reed-Berendt, B. G.; Melen, R. L.; Morrill, L. C. FLP-Catalyzed Transfer Hydrogenation of Silyl Enol Ethers. *Angew. Chem., Int. Ed.* **2018**, *57*, 12356–12359. (c) Webb, J. D.; Laberge, V. S.; Geier, S. J.; Stephan, D. W.; Crudden, C. M. Borohydrides from Organic Hydrides: Reactions of Hantzsch's Esters with $B(C_6F_5)_3$. *Chem. - Eur. J.* **2010**, *16*, 4895–4902.

(23) For reviews, see: (a) Irrgang, T.; Kempe, R. 3d-Metal Catalyzed N- and C-Alkylation Reactions via Borrowing Hydrogen or Hydrogen Autotransfer. *Chem. Rev.* **2019**, *119*, 2524–2549. (b) Reed-Berendt, B. G.; Polidano, K.; Morrill, L. C. Recent Advances in Homogeneous Borrowing Hydrogen Catalysis Using Earth-Abundant First Row Transition Metals. *Org. Biomol. Chem.* **2019**, *17*, 1595–1607.

(24) See the [Supporting Information](#) for further details of optimization. Electron-rich diaryl amines are expected to be better hydride donors and, thus, undergo the $B(C_6F_5)_3$ C–H cleavage more readily. For studies of C–H hydride donor abilities, see: Ilic, S.; Alherz, A.; Musgrave, C. B.; Glusac, K. D. Thermodynamic and Kinetic Hydricities of Metal-Free Hydrides. *Chem. Soc. Rev.* **2018**, *47*, 2809–2836.

(25) Amine **6a** was found to be a more-general methylating agent versus **4a**.

(26) The addition of TMP significantly improved the yield when NH indoles were used. See the [Supporting Information](#).

(27) Guidotti, S.; Camurati, I.; Focante, F.; Angellini, L.; Moscardi, G.; Resconi, L.; Leardini, R.; Nanni, D.; Mercandelli, P.; Sironi, A.; Beringhelli, T.; Maggioni, D. Synthesis and Reactivity of $(C_6F_5)_3B$ -N-Heterocycle Complexes. 1. Generation of Highly Acidic sp^3 Carbons in Pyrroles and Indoles. *J. Org. Chem.* **2003**, *68*, 5445–5465. (b) Focante, F.; Camurati, I.; Nanni, D.; Leardini, R.; Resconi, L. Synthesis and Reactivity of N-Heterocycle- $B(C_6F_5)_3$ Complexes. 3. Generation of N-Methylpyrrol-2-yl and N-Methylindol-2-yl Borate Zwitterions with Acidic sp^3 Carbons. *Organometallics* **2004**, *23*, 5135–5141.

(28) For an example where 3,3'-bisindolylmethane products form, see ref **7**.

(29) For an example of the impressive functional group compatibility of catalytic $B(C_6F_5)_3$ processes, see: Bender, T. A.; Payne, P. R.; Gagné, M. R. Late-Stage Chemoselective Functional-Group Manipulation of Bioactive Natural Products with Super-Electrophilic Silylium ions. *Nat. Chem.* **2018**, *10*, 85–90.

(30) The amine byproduct (cf. amine **18**, Scheme 6) was recovered in 98% yield.

(31) Benzylating agent **4b** was marginally better over the analogous alkylating agent based on amine **6**.

(32) Benzofurans, furans, benzothiophenes, and thiophenes were unreactive, and pyrroles gave low yields. See the [Supporting Information](#) for details.

(33) Schneider, A. F.; Chen, Y.; Brook, M. A. Trace Water Affects Tris(pentafluorophenyl)borane-Catalytic Activity in the Piers–Rubinsztajn Reaction. *Dalton Trans.* **2019**, *48*, 13599–13606.

(34) For example, see: (a) Thomson, J. W.; Hatnean, J. A.; Hastie, J. J.; Pasternak, A.; Stephan, D. W.; Chase, P. A. Improving the Industrial Feasibility of Metal-Free Hydrogenation Catalysts Using Chemical Scavengers. *Org. Process Res. Dev.* **2013**, *17*, 1287–1292.

(35) In the case of oxindoles **9a**, yields were improved upon removal of the $O(SiEt_3)_2$ byproduct by simply applying a vacuum. See the [Supporting Information](#).

(36) (a) Schwartz, T. L.; Siddiqui, U. A.; Stahl, S. M. Vilazodone: A Brief Pharmacological and Clinical Review of the Novel Serotonin Partial Agonist and Reuptake Inhibitor. *Ther. Adv. Psychopharmacol.* **2011**, *1*, 81–87. (b) Gründer, G.; Wetzel, H.; Hammes, E.; Benkert, O. Roxindole, a Dopamine Autoreceptor Agonist, in the Treatment of Major Depression. *Psychopharmacology* **1993**, *111*, 123–126. (c) Sanchez, C.; Papp, M. The Selective Sigma2 Ligand Lu 28–179 has an Antidepressant-Like Profile in the Rat Chronic Mild Stress Model of Depression. *Behav. Pharmacol.* **2000**, *11*, 117–124. (d) Marchese, G.; Rui, S.; Casti, P.; Bartholini, F.; Saba, P.; Gessa, G. L.; Pani, L. Carmoxirole is Able to Reduce Amisulpride-Induced Hyperprolactinemia Without Affecting its Central Effect. *Eur. J. Pharmacol.* **2002**, *447*, 109–114.

(37) Wasa has previously shown that *N*-aryl pyrrolidines undergo hydride abstraction; see ref **19**.

(38) Electron-rich aromatic rings can be removed from nitrogen under oxidative conditions (cf. PMP protecting groups). For a relevant example, see ref **19b**.

(39) Partial deuterium incorporation at the *N*-CH₃ site of **2a-d₃** may indicate that hydride abstraction may also occur at *N*-CH₃, or that a 1,5-prototropic shift in intermediate **19** (*R* = CH₃) occurs prior to hydride transfer (cf. [Scheme 6b](#), step iv).

(40) Attempts to prevent the elimination and isolate the corresponding aminomethylation derivative of **17** were unsuccessful. For a related elimination-reduction sequence, see: Deb, M. L.; Baruah, P. K. Deamination of Indole Mannich Bases: An Efficient Route to 3-Benzyl/Alkylindoles via a Metal-Free Transfer Hydrogenation Under Microwave Irradiation. *COCAT* **2015**, *3*, 84–89.